TYLENOL (acetaminophen)

DO NOT GIVE ACETAMINOPHEN TO BABIES UNDER 3 MONTHS OLD without calling your doctor first.
ALWAYS READ LABEL FIRST, since different medications have different concentrations.
ALWAYS USE MEASURING DEVICE (dropper, syringe or measuring cup) that comes with the medication or you could risk overdosing. Never use droppers from different medications or spoons from the kitchen.

		Infant's Drops 80mg/0.8ml	*Infant's Suspension 160 mg per 5 mL *1 tsp = 5 ml	Children's Liquid 160mg/tsp	Soft chews And Chewable tablets 80mg/each	Junior Chewable Tablets 160mg	Fever-all Suppository 120,325 & 650 mg	Adult 325 mg
Dose by Weight	**Age**	**Dropper**	**Syringe**	**Teaspoon**	**Tablet**	**Tablet**	**Suppository**	**Tablet**
6-11 lbs	0-3mos	1/2 (0.4ml)	1/4syringe (1.25 ml)				1/3 of 120 mg	
12-17lbs	4-11 mos	1 (0.8ml)	1/2 syringe (2.5mL)	1/2 tsp.			2/3 of 120 mg	
18-23lbs	12-23 mos	1 -1/2 (1.2ml)	3/4syringe (3.75 ml)	3/4 tsp.			120 mg	
24-35lbs	2-3 years	2 (1.6 ml)	1 syringe (5 mL)	1 tsp.	2 tablets		1-1/2 120mg **or** 1/2 325 mg	
36-47lbs	4-5 years			1 -1/2 tsp.	3 tablets	1 ½ tablets	3/4 325 mg	
48-59lbs	6-8 years			2 tsp.	4 tablets	2 tablets	325 mg	1 tablet
60-71lbs	9-10 years			2 1/2 tsp.	5 tablets	2 ½ tablets	3/4 650 mg	
72-95lbs	11 years			3 tsp.	6 tablets	3 tablets	3/4 650 mg	
96-119lbs	12 years					4 tablets	650 mg	2 tablets
120lbs								2 tablets
150lbs								3 tablets